

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

Table 1. Selected Electronic State/Valence Distributions^a

Elm	Chrg	Rad	Bd#1	Bd#2	#Extern Bds ^b	#e's Contrib	Shorthand	Structure
B			1	2	0	1	[120/1]	B-1
B			1	1	1	0	[111/0]	B-2
B	-1		1	2	1	1	[121/1-]	B-3
C			1	2	1	1	[121/1]	C-1
C	+1		1	1	1	0	[111/0+]	C-2
C	-1		1	1	1	2	[111/2-]	C-3
C		•	1	1	1	1	[111/1*]	C-4
C			1	1	2	1	[112/1]	C-5
N			1	2	0	1	[120/1]	N-1
N			1	1	1	2	[111/2]	N-2
N	+1		1	2	1	1	[121/1+]	N-3
N	+1	•	1	1	1	1	[111/1+*]	N-4
O			1	1	0	2	[110/2]	O-1
O	+1		1	1	1	2	[111/2+]	O-2
O	+1		1	2	0	1	[120/1+]	O-3
P like N ^c								
P			1	2	2	1	[122/1]	P-1
S like O								
S			2	2	0	2	[220/2]	S-1
Cl	+1		1	1	0	2	[110/2+]	Cl-1
Cl			1	2	2	1	[122/1]	Cl-2
Cl			1	2	4	1	[124/1]	Cl-3

FIG. 7




FIG. 8

Table 2. Procedure Control Flags

Flag	Meaning
kDontAssumeImpH	Otherwise, heteroatoms might carry undrawn hydrogens.
kIfFailWithOneSystem_FailWithAll	If given two or more delocalized systems and one fails, the molecule is returned unchanged, and the procedure fails.
kDoNotCreateCharges	Do not create zwitterions, i.e., more charges than necessary to achieve the system's net charge.
kDoNotCreateRadicals	Do not create more than one radical.
kConfineChargesToHeteroatoms	All charged atoms must be heteroatoms.
kConfineRadicalsToHeteroatoms	All atoms with an unpaired electron must be heteroatoms.
kFavorMultiplyBondedHetero	When a system can support more or fewer multiple bonds, favor the form with more multiple bonds (even if it is anti-aromatic).
kDisfavorAntiaromaticSystems	Use this flag in conjunction with the previous.
kSolutionMustBeFullyAlternating	Bonds must alternate as single and double.




FIG. 10

Table 3. Meanings of bits in the ESVD Screening Bitmask (ESB) and Atom Screening Bitmask (ASB)

<u>Bit #</u>	<u>Description</u>	<u>Bit #</u>	<u>Description</u>
0	Has an internal single bond	8	Charge = 0
1	Has two internal single bonds	9	Charge = +1
2	Has an internal double bond	10	Charge = -1
3	Has two internal double bonds	11	Charge ≠ -1
4	Has an external bond	12	Charge ≠ +1
5	Does <i>not</i> have an external bond	13	Radical present
		14	Radical not present

FIG. 11




FIG. 12

<u>Action #</u>	<u>Action</u>	<u>Atom / Bond</u>	<u>Action #</u>	<u>Action</u>	<u>Atom or Bond</u>
1	Assign ESVD	Atom 0	14	Assign Bond	Bond 10
2	Assign Bond	Bond 0	15	Assign ESVD	Atom 9
3	Assign ESVD	Atom 1	16	Assign Bond	Bond 9
4	Assign Bond	Bond 1	17	Assign ESVD	Atom 8
5	Assign ESVD	Atom 2	18	Assign Bond	Bond 8
6	Assign Bond	Bond 2	19	Assign ESVD	Atom 7
7	Assign ESVD	Atom 3	20	Assign Bond	Bond 7
8	Assign Bond	Bond 3	21	Assign ESVD	Atom 6
9	Assign ESVD	Atom 4	22	Assign Bond	Bond 6
10	Assign Bond	Bond 4	23	Verify completed	Atom 4
11	Assign ESVD	Atom 5	24	Complete	---
12	Assign Bond	Bond 5			
13	Verify completed	Atom 0			

FIG. 13

Table 4. The Actions Comprising a Strategy

<u>Action</u>	<u>Applies To</u>	<u>Explanation</u>
Assign ESVD	Atom	Find the ESVD's for the current atom that are compatible with its environment. The best one is used directly, and if there is more than one, the rest are queued.
Assign Bond	Bond	Assign a bond order to the current bond, consistent with the ESVD of the previous atom, i.e. the (earliest occurring) atom adjacent to the bond. (The bond's other atom has not been encountered yet, unless the bond closes a ring. Even in this case, the other atom's environment is not taken into account. It will be checked in the next Action.)
Verify Completed	Atom	This Action is taken just after the last bond in a ring or acyclic chain is fixed. Ordinarily the bonds of an atom are sure to be compatible with its assigned ESVD because its ESVD was picked to be compatible with the bond leading to it, and the bond leading away from it was selected to be compatible with its ESVD. However, a ring closure atom has not had its ESVD checked with respect to the ring closure bond, nor has a terminal atom in an acyclic chain. Thus, in this Action the atom is checked to verify that its final bonding environment is compatible with its ESVD.
Complete	---	Signifies that the path is completed, and all atoms have been assigned compatible ESVD's and bond orders. If the net charge or radical count of the putative solution is wrong, the solution is rejected. If the solution is perfect, as defined elsewhere, it is returned directly and the procedure terminates. Otherwise, if it is the best solution yet, it displaces the previous best candidate.




FIG. 15

FIG. 16






FIG. 17




FIG. 18




FIG. 19




FIG. 20


```

graph TD
    1010[Select an initial atom (step 1010).] --> 1015[Assign a bond order to an adjacent bond (step 1015)]
    1015 --> 1020[Assign, to a next adjacent atom, an ESVD that is consistent with the previous bond (step 1020)]
    1020 --> 1030[Select the next adjacent bond's order (step 1030)]
    1030 --> 1040[Repeat the same approach for additional atoms (step 1040)]
    1040 --> 1050{If an unacceptable state is encountered}
    1050 --> 1055[Reject the path values (step 1050)]
    1055 --> 1060[Backtrack to the last point where a selection was made and proceed forward from there with a different selection (step 1060)]
    1060 --> 1070{If a solution is found that cannot be improved upon, terminate (step 1070)}
    1070 --> 1080[If a solution is found that is not optimal and there are other choices, explore the other choices (step 1080)]

```

FIG. 22

[illegible]

```

graph TD
    A[Note the total charge and radical count required of the pi system (step 2070)] --> B[Calculate the internal coordination number ("ICN") of each atom (step 2080)]
    B --> C[Identify bonds in DS that must be fixed, as a consequence of having an adjacent fixed multiple bond (step 2090)]
    C --> D[Develop a path through DS (step 2100)]
    D --> E[Construct the strategy list (step 2110)]
    E --> F[Identify ring systems (step 2120)]
  
```



A diagram showing a block labeled 'E' hanging from a horizontal line representing a string. An arrow points downwards from the string to the block, indicating the force of gravity.

FIG. 23A




FIG. 23B

3000
↓




FIG. 24

00506747 021800

45322
 45323
 45324
 45325
 45326
 45327
 45328
 45329
 45330
 45331
 45332
 45333
 45334
 45335
 45336
 45337
 45338
 45339
 45340
 45341
 45342
 45343
 45344
 45345
 45346
 45347
 45348
 45349
 45350
 45351
 45352
 45353
 45354
 45355
 45356
 45357
 45358
 45359
 45360
 45361
 45362
 45363
 45364
 45365
 45366
 45367
 45368
 45369
 45370
 45371
 45372
 45373
 45374
 45375
 45376
 45377
 45378
 45379
 45380
 45381
 45382
 45383
 45384
 45385
 45386
 45387
 45388
 45389
 45390
 45391
 45392
 45393
 45394
 45395
 45396
 45397
 45398
 45399
 45400




Fig. 25A




FIG. 25B




FIG. 25C

00506717-004000

5000




FIG. 26A




FIG. 26B

6000
→




FIG 27

00506717-024900

7000 →




FIG. 28

005733 2 299560

8000




FIG. 29

00506717-001000

9000
↓

Initialize ASB to zero (step 9010)

Set bits of ASB according to the number of internal bonds adjacent to *a* that are fixed, with bond order single (step 9020)

Set bits according to the number of internal bonds adjacent to *a* that are fixed, with bond order double (step 9030)

If *a* has more than two adjacent bonds, set bit {4} (step 9040)

If *a* has exactly two adjacent bonds, and if the control flags dictate that no implicit hydrogens exist, and *a*'s element type is not carbon, set bit {5} (step 9050)

If the residual radical value of *S* is "no-radical", and the control flags do not permit unnecessary creation of radicals, set bit {14} (step 9060)

If the residual charge of *S* is zero and the control flags do not permit unnecessary creation of charged atoms; or if *a*'s element type is carbon and the control flags require charges to be situated on heteroatoms, set bits {11,12} (step 9070)

Otherwise

If the residual charge of *S* is positive, set bit {11} (step 9075)

If the residual charge of *S* is negative, set bit {12} (step 9080)

Return ASB (step 9090)

FIG. 30

For each ring r belonging to the RB, execute steps 10050 - 10110 (step 10040)

2.	
----	--

7

If e is odd, subtract one from R (step 10080)

↓

5

Adjust R by the one-ring penalty amount, where r_2 is substituted for r (step 10110)

FIG. 31A

[illegible]

If there are three or more rings in RB

Take $r3$ to be the compound ring represented by P
(step 10120)

Adjust R by the one-ring penalty amount, where $r3$ is
substituted for r (step 10130)

Apply a significant penalty for gratuitous charges and radicals
(steps 10160 - 10180) (step 10140)

Take NSC to be the number of superfluous charges (step
10160)

Take NSR to be the number of superfluous radicals (step
10170)

Reduce the score by ten times $(NSC + NSR)$ (step 10180)

Apply a small penalty for locating a charge on a carbon rather than
an available heteroatom (steps 10200 - 10220) (step 10190)

Take NCC to be the number of charged carbon atoms in S
(step 10200)

Take NNH to be the number of neutral heteroatoms in S
(step 10210)

If NCC exceeds NNH , reduce R by $(NCC - NNH)$ (step 10220)

Penalize pairs of adjacent atoms having identical, either empty or
full, orbitals (step 10230)

D

FIG. 31B




FIG. 31C

09506717-021800